

CONCLUSIONS

In conclusion, the UHPLC-MS method with ultrafiltration described in this study demonstrated a good capability for determining unbound VPA concentrations. The method has advantages of easy sample preparation and rapid analysis, and it can be used to facilitate large-scale investigations of the relationship between unbound VPA concentrations and clinical outcomes in epileptic children.

REFERENCES

- Gibbs HG, Zimmerman DE, Shermock KM, et al. Comparison of free fraction serum valproic acid concentrations between inpatients and outpatients. *Am J Health Syst Pharm*. 2015;72:121–126.
- Ferraro TN, Buono RJ. The relationship between the pharmacology of antiepileptic drugs and human gene variation: an overview. *Epilepsy Behav*. 2005;7:18–36.
- Zaccara G, Messori A, Moroni F. Clinical pharmacokinetics of valproic acid. *Clin Pharmacokinet*. 1988;15:367–389.
- Rapeport WG, Mendelow AD, French G, et al. Plasma protein-binding and CSF concentrations of valproic acid in man following acute oral dosing. *Br J Clin Pharmacol*. 1983;16:365–369.
- Itoh H, Suzuki Y, Fujisaki K, et al. Correlation between plasma ammonia level and serum trough concentration of free valproic acid in patients with epilepsy. *Biol Pharm Bull*. 2012;35:971–974.
- Suzuki Y, Itoh H, Abe T, et al. No effect of co-administered antiepileptic drugs on in-vivo protein binding parameters of valproic acid in patients with epilepsy. *J Pharm Pharmacol*. 2011;63:976–981.
- Zhang ZQ, Dong WC, Yang XL, et al. The influence of plasma albumin concentration on the analysis methodology of free valproic acid by ultrafiltration and its application to therapeutic drug monitoring. *Ther Drug Monit*. 2015;15:361–365.
- Dwivedi R, Gupta YK, Singh M, et al. Correlation of saliva and serum free valproic acid concentrations in persons with epilepsy. *Seizure*. 2015; 25:187–190.
- March C, Blanke RV. Determination of free valproic acid concentrations using the Amicon Micropartition MPS1 ultrafiltration system. *Ther Drug Monit*. 1985;7:115–120.
- Gao S, Miao H, Tao X, et al. LC-MS/MS method for simultaneous determination of valproic acid and major metabolites in human plasma. *J Chromatogr B Analyt Technol Biomed Life Sci*. 2011;879:1939–1944.
- Zhao M, Li G, Qiu F, et al. Development and validation of a simple and rapid UPLC-MS assay for valproic acid and its Comparison with immunoassay and HPLC methods. *Ther Drug Monit*. 2016;38:246–252.
- FDA. *Guidance for Industry Bioanalytical Method Validation [web site]*. 2013. Available at: <https://www.fda.gov/downloads/drugs/guidancecomplianceregulatoryinformation/guidances/ucm368107.pdf>. Accessed March 15, 2017.
- Matuszewski BK, Constanzer ML, Chavez-Eng CM. Strategies for the assessment of matrix effect in quantitative bioanalytical methods based on HPLC-MS/MS. *Anal Chem*. 2003;75:3019–3030.
- Riva R, Albani F, Baruzzi A, et al. Determination of unbound valproic acid concentration in plasma by equilibrium dialysis and gas-liquid chromatography: methodological aspects and observations in epileptic patients. *Ther Drug Monit*. 1982;4:341–352.
- Riva R, Albani F, Cortelli P, et al. Diurnal fluctuations in free and total plasma concentrations of valproic acid at steady state in epileptic patients. *Ther Drug Monit*. 1983;5:191–196.
- Cheng H, Liu Z, Blum W, et al. Quantification of valproic acid and its metabolite 2-propyl-4-pentenoic acid in human plasma using HPLC-MS/MS. *J Chromatogr B Analyt Technol Biomed Life Sci*. 2007;850: 206–212.
- Ueshima S, Aiba T, Makita T, et al. Characterization of non-linear relationship between total and unbound serum concentrations of valproic acid in epileptic children. *J Clin Pharm Ther*. 2008;33:31–38.

ERRATUM

Prospective Randomized Trial Investigating the Influence of Pharmaceutical Care on the Intra-Individual Variability of Tacrolimus Concentrations Early After Kidney Transplant: Erratum

In the August 2016 issue of Therapeutic Drug Monitoring in the article “Prospective Randomized Trial Investigating the Influence of Pharmaceutical Care on the Intra-Individual Variability of Tacrolimus Concentrations Early After Kidney Transplant”, the author Juliana F. Mansur should be listed as Juliana B. Mansur.

REFERENCE

- Bessa AB, Felipe CR, Hannun P, et al. Prospective randomized trial investigating the influence of pharmaceutical care on the intra-individual variability of tacrolimus concentrations early after kidney transplant. *Ther Drug Monit*. 2016;38:447–455.